

# Polar Bioactive Constituents from Aerial Parts of *Thymus longicaulis* C. Presl

Brigida D'Abrosca<sup>1\*</sup>, Monica Scognamiglio<sup>2</sup>, Vittoria Graziani<sup>1</sup>, Antonio Fiorentino<sup>1</sup>

<sup>1</sup>Department of Environmental Biological and Pharmaceutical Sciences and Technologies, Second University of Naples, Italy

<sup>2</sup>Max Planck Institute for Chemical Ecology, Hans-Knöll-Straße, Germany

**Corresponding authors:** Brigida, D'Abrosca. Department of Environmental Biological and Pharmaceutical Sciences and Technologies, Second University of Naples, Via Vivaldi 43, Caserta, Italy; Tel: +39-0823-274564; E-mail: [brigida.dabrosca@unina2.it](mailto:brigida.dabrosca@unina2.it)

## Abstract

*Thymus longicaulis* C. Presl., belonging to the Lamiaceae family, is a small aromatic perennial herb typical of the Mediterranean vegetation. This species, known in traditional phytoterapy of Italy, has been extensively investigated in terms of chemical analysis and biological activity of its essential oils. Nevertheless, few data are available in the literature, regarding the chemical characterization of polar components of *T. longicaulis*. In this study, the phytochemical investigation of methanol extract of *T. longicaulis* through different chromatographic techniques, led to the isolation of thirteen compounds. The structures of rosmarinic acid and two derivatives, as well as that of flavones, triterpenes and a lignan have been elucidated on the basis of extensive NMR spectroscopic analyses. The evaluation of DPPH radical scavenging activity of pure compounds has been performed.

**Received Date: 22 February, 2016**

**Accepted Date: 29 February, 2016**

**Published Date: 3 March, 2016**

**Citation:** D'Abrosca, B., et al. Polar Bioactive Constituents from Aerial Parts of *Thymus Longicaulis* C. Presl. (2016) Lett Health Biol Sci 1(1): 1- 4.

**DOI:** 10.15436/2475-6245.16.002



## Introduction

Plants are a rich source of bioactive compounds, characterized by a wide range of pharmacological activities, can be used for different applications such as health promoting ingredients, nutraceuticals, and food additives in formulations of functional foods<sup>[1]</sup>. Although fresh and dried aromatic plants, due to their richness in volatile components, have been used as flavorings since ancient times, they produce a large amount of other secondary metabolites (e.g. flavonoids, phenolic acids, saponins), responsible for several beneficial effects on human health. By virtue of their properties, during the last few decades, they have also become a subject for a search of natural antioxidants<sup>[2]</sup>. In particular, flavonoids and phenolic acids play an important role in protecting organisms against dangerous effects of reactive oxygen species (ROS). So recently, there has been a great interest towards the therapeutic potentials of aromatic plants as antioxidant in reducing free radical induced tissue damage and their use as antioxidant in food and drugs<sup>[3]</sup>.

The genus *Thymus*, belonging to the Lamiaceae family, is one of the most critical genera of the Mediterranean flora. This genus comprises about 400 species of perennial aromatic herbaceous plants one of them is *Thymus longicaulis* (C. Presl). *T. longicaulis* is a species with long, somewhat woody, creeping branches, with a terminal inflorescence<sup>[4]</sup>. This species, known in traditional phytoterapy of South and Central Italy, is used as a tonic as against cough and influence<sup>[5]</sup>.

Recently, chemical composition and different biological properties of essential oils of *T. longicaulis* have been investigated: antimicrobial activity of *T. longicaulis* essential oil from Croatia<sup>[6]</sup>, antioxidant properties of essential oil of *T. longicaulis* from Turkey<sup>[7]</sup>.

Few reports<sup>[8]</sup> are available regarding the chemical characterization of polyphenolic components of *T. longicaulis*. In the investigation of in this framework, the aim of this study is the phytochemical investigation of the polar extract of *T. longicaulis* as well as the evaluation of DPPH radical scavenging activity of the isolated compounds.

## Materials and Methods

### General experiment procedures

NMR spectra were recorded at 300.03 MHz for <sup>1</sup>H and 75.45 MHz for <sup>13</sup>C on a Varian Mercury 300 spectrometer Fourier transform NMR in CD<sub>3</sub>OD or CDCl<sub>3</sub> solutions at 25°C. Chemical shifts are reported in δ (ppm) and referenced to the residual sol-

vent signal,  $J$  (coupling constant) are given in Hz. Standard pulse sequences and phase cycling from Varian library were used for  $^1\text{H}$ ,  $^{13}\text{C}$ , DEPT, DQF-COSY, COSY, TOCSY, HSQC, H2BC, HMBC and CIGAR-HMBC experiments.  $^1\text{H}$  NMR spectra were acquired over a spectral window from 14 to 2 ppm, with 1.0 s relaxation delay, 1.70 s acquisition time (AQ),  $90^\circ$  pulse width = 13.8  $\mu\text{s}$ . The initial matrix was zero-filled to 64 K.  $^{13}\text{C}$ -NMR spectra were recorded in  $^1\text{H}$  broadband decoupling mode, over a spectral window from 235 to 15 ppm, 1.5 s relaxation delay,  $90^\circ$  pulse width = 9.50  $\mu\text{s}$ , AQ = 0.9 s. The number of scans for both  $^1\text{H}$  and  $^{13}\text{C}$ -NMR experiments was chosen depending on the concentration of the samples. Also for homonuclear and heteronuclear 2D-NMR experiments, data points, number of scan and of increments were adjusted according to the sample concentrations. Correlation spectroscopy (COSY) and double quantum filtered COSY (DQF-COSY) spectra were recorded with gradient enhanced sequence at spectral widths of 3000 Hz in both  $f_2$  and  $f_1$  domains; the relaxation delays were of 1.0s. The total correlation spectroscopy (TOCSY) experiments were performed in the phase-sensitive mode with a mixing time of 90 ms. The spectral width was 3000 Hz. Nuclear Overhauser effect spectroscopy (NOESY) experiments were performed in the phase-sensitive mode. The mixing time was 500 ms and the spectral width was 3000 Hz. For all the homonuclear experiments, the initial matrix of 512 x 512 data points was zero-filled to give a final matrix of 1 k x 1 k points. Proton- detected heteronuclear correlations were measured.

Heteronuclear single-quantum coherence (HSQC) experiments (optimized for  $^1J(\text{H,C}) = 140$  Hz) were performed in the phase sensitive mode with field gradient; the spectral width was 12,000 Hz in  $f_1$  ( $^{13}\text{C}$ ) and 3000 Hz in  $f_2$  ( $^1\text{H}$ ) and 1.0 s of relaxation delay; the matrix of 1 k x 1 k data points was zero-filled to give a final matrix of 2 k x 2 k points. Heteronuclear 2 bond correlation (H2BC) spectra were obtained with  $T = 30.0$  ms, and a relaxation delay of 1.0 s; the third-order low-pass filter was set for  $130 < ^1J(\text{C,H}) < 165$  Hz. Heteronuclear multiple bond coherence (HMBC) experiment (optimized for  $nJ(\text{H,C}) = 8$  Hz) was performed in the absolute value mode with field gradient; typically,  $^1\text{H}$ - $^{13}\text{C}$  gHMBC were acquired with spectral width of 18,000 Hz in  $f_1$  ( $^{13}\text{C}$ ) and 3000 Hz in  $f_2$  ( $^1\text{H}$ ) and 1.0 s of relaxation delay; the matrix of 1 k x 1 k data points was zero-filled to give a final matrix of 4 k x 4 k points. Constant time inverse-detection gradient accordion rescaled heteronuclear multiple bond correlation spectroscopy (CIGAR-HMBC) spectra ( $8 > nJ(\text{H,C}) > 5$ ) were acquired with the same spectral width used for HMBC. Heteronuclear single-quantum coherence-total correlation spectroscopy (HSQC-TOCSY) experiments were optimized for  $nJ(\text{H,C}) = 8$  Hz, with a mixing time of 90 ms.

Analytical TLC was performed on Merck Kieselgel 60  $F_{254}$  or RP-8  $F_{254}$  plates with 0.2 mm layer thickness. Spots were visualized by UV light or by spraying with  $\text{H}_2\text{SO}_4/\text{AcOH}/\text{H}_2\text{O}$  (1:20:4). The plates were then heated for 5 min at  $110^\circ\text{C}$ . Preparative TLC was performed on Merck Kieselgel 60  $F_{254}$  plates, with 0.5 or 1.0 mm film thickness. Column chromatography (CC) was performed on Merck Kieselgel 60 (70–240  $\mu\text{m}$ ), Merck Kieselgel 60 (40–63  $\mu\text{m}$ ), Bakerbond C8 and C18 Sephadex LH-20, Amberlite XAD-4.

## Plant material

*Thymus longicaulis* C. Presl was collected in a garrigue on the calcareous hills of Durazzano, ( $41^\circ 3' \text{N}$ ,  $14^\circ 27' \text{E}$ ; southern Italy) in the vegetative state and identified by Dr. Assunta Esposito of the Dept. of Environmental, Biological and Pharmaceutical Sciences and Technologies of Second University of Naples (SUN). A voucher specimen (CE235) has been deposited at the Herbarium of the Department. Leaves of *Thymus longicaulis* were harvested and immediately frozen in liquid  $\text{N}_2$  in order to avoid unwanted enzymatic reactions and stored at  $-80^\circ\text{C}$  up to the freeze drying process. Once freeze dried they were powdered in liquid nitrogen and stored at  $-20^\circ\text{C}$  until the extraction process was carried out.

## Extraction and isolation of compounds

Dried leaf material was powdered and extracted by ultrasound assisted extraction (Elma<sup>®</sup> Transonic Digital) one hour with methanol. The extract was filtered on Whatman paper and concentrated under *vacuum*. After removal of the solvent, a dried crude extract was obtained (4.1 g) which was stored at  $-20^\circ\text{C}$  until its purification. The methanol extract, dissolved in distilled water and shaken with EtOAc, give an aqueous and an organic fraction.

The first was chromatographed on Amberlite XAD-4 and eluted first with water, to eliminate sugars, peptides, free amino acids and other primary metabolites, and then with methanol. The alcoholic eluate furnished 1.1g of residual material which was chromatographed on Sephadex LH-20 eluting with MeOH/ $\text{H}_2\text{O}$  polarity decreasing solutions and collecting six fractions (A-F). Fraction A, re-chromatographed by  $\text{SiO}_2$ -flash CC, eluting with the organic phase of a biphasic solution constituted by  $\text{CHCl}_3/\text{MeOH}/\text{H}_2\text{O}$  (13:7:4), furnished pure compound **1** (7.7 mg) and another fraction that, purified by TLC [ $\text{SiO}_2$ , lower phase of  $\text{CHCl}_3/\text{MeOH}/\text{H}_2\text{O}$  (13:7:4)] gave compounds **2** (1.2 mg) and **13** (1.6 mg).

Fraction B, was chromatographed by RP-18 CC furnishing two fractions: fraction B1 contained pure **6** (20.1 mg), while B2 gave pure compound **7** (4.5 mg). Fraction C contained pure compound **4** (89 mg), fraction D, instead, re-chromatographed by RP-18 CC furnished a fraction identified as compound **9** (10.9 mg). Fraction E was purified by  $\text{SiO}_2$  TLC (0.5 mm), eluting with the lower phase of the biphasic solution  $\text{CHCl}_3/\text{MeOH}/0,1\%$  TFA (13:7:2), and gave two spots. The first spot was identified as pure **11** (2.8 mg), while the second spot as the metabolite **10** (1.5 mg). Finally, fraction F after re-chromatography on RP-18 CC (MeOH:  $\text{H}_2\text{O}$  polarity decreasing solutions) furnished a fraction identified as compound **12** (4.5 mg).

The organic fraction of methanol extract, chromatographed on Sephadex LH-20 eluting with hexane/MeOH/ $\text{CHCl}_3$  (2:1:1) solution, furnished three fractions G1-G3. The first re-chromatographed by  $\text{SiO}_2$ -flash CC, eluting with decreasing polarity of  $\text{CHCl}_3/\text{MeOH}$  solutions, furnished pure compound **4** (33.3 mg).


Fraction G2, purified by  $\text{SiO}_2$ -flash CC, eluting with decreasing polarity of  $\text{CHCl}_3/\text{EtOAc}$  solution, gave a fraction identified as compound **3** (6.6 mg). Finally, fraction G3 was purified by TLC,  $\text{SiO}_2$   $\text{CHCl}_3/\text{MeOH}$  (4:1) to obtain pure **8** (1.2 mg).

### Radical scavenging capacity


In order to assess the antioxidant efficacy of the isolated pure metabolites from *T. longicaulis* leaves, the 2, 2-diphenyl-1-picrylhydrazyl (DPPH) spectrophotometric method was performed as previously reported<sup>[9]</sup>. Increasing concentrations of pure metabolites were tested (5, 25, 50, 100 and 250  $\mu$ M). All the tests and analyses were carried out in triplicate. Trolox<sup>®</sup> (6-hydroxy- 2, 5, 7, 8-tetramethylchroman-2-carboxylic acid), a water soluble vitamin E analogue, was used as a positive control.

### Results and Discussion

The phytochemical investigation of the methanol extracts of thyme led to isolation and characterization of thirty compounds (figures 1 and 2) belonging to different classes of secondary metabolites: isoprenoids, cinnamic acid derivatives, and flavones. In particular compounds **1** and **2** were identified as tuberonic acid 13-O- $\beta$ -D-glucopyranoside and 4, 7-megastigma-dien-3-one 9-O- $\beta$ -D-glucopyranoside, respectively, both isolated from seeds of *Astragalus complanatus*<sup>[10]</sup>.



**Figure 1:** Chemical structures of terpenoids (1- 4) isolated from *Thymus longicaulis*.



**Figure 2:** Chemical structures of caffeic acid derivatives (5-8), flavones (9-12), and lignan (13) isolated from *Thymus longicaulis*.

Metabolites **3** and **4** are two ursane triterpenes. In particular, **3** was identified as 3 $\beta$ -hydroxy-urs-12-en-28-oic acid, known as ursolic acid, while compound **4** has been character-

ized as pomolic acid, by comparison of its spectral data with those reported in literature data. Both triterpenes were as reported as constituent of *Annurca apple* fruits<sup>[11]</sup>. Metabolites **5-7** were identified as caffeic acid derivatives (figure 2); in particular, compound **5** was recently reported as constituent of several aromatic Mediterranean plant species<sup>[8]</sup>. Compound **6** was elucidated as salvianolic acid K, previously isolated from *Salvia deserta*<sup>[12]</sup>.


NMR data of **7**, known as 4-O- $\beta$ -D-glucopyranosyl rosmarinic acid, was in good accordance with<sup>[13]</sup>, that reported this compound as constituent of *Sanicula lamelligera* while metabolite **8** was identified as methyl caffeoate.

Compounds **9-12** were identified as flavone derivatives (figure 2). Compound **9** was previously isolated from aerial parts of *Teucrium polium*<sup>[14]</sup>, while luteolin-7-O- $\beta$ -D-glucopyranoside, known as cynaroside (**10**), was already reported as constituent of *Marrubium globosum ssp. libanoticum*<sup>[15]</sup>.

Flavone **11** was characterized as isoscutellarein 7-O- $\beta$ -D-glucopyranoside, by comparison of its spectral data with those reported in literature data<sup>[16]</sup>. Compound **12** was identified as gossipytrin first isolated from *Papaver nudicaule*<sup>[17]</sup>.

Finally, compound **13** identified as (+)-pinoselinol 4, 4'-O-bis- $\beta$ -D-glucopyranoside was reported from constituent of *Clematis stans* roots<sup>[18]</sup>.

In order to evaluate the antioxidant efficacy of the isolated pure metabolites from *T. longicaulis* leaves, the 2, 2-diphenyl-1-picrylhydrazyl (DPPH) spectrophotometric method was performed as previously reported using five decreasing concentrations of pure metabolites. The results are reported in (figure 3). The polyphenols showed the highest radical scavenging activity. In particular rosmarinic acid (**5**) and salvianolic acid (**6**) reducing DPPH concentration of 94 % and 67 %, respectively at highest concentration, with considerable radical scavenging activity also at 100 $\mu$ M.



**Figure 3:** DPPH radical scavenging activity of compounds isolated from *Thymus longicaulis*.

Rosmarinic acid is distributed in 26 plant families, and its biological activity has been extensively examined. Earlier studies ascribed to rosmarinic acid antiviral, antibacterial, anti-inflammaroty and antioxidant properties<sup>[19]</sup>. Recently, others interesting biological effects of rosmarinic acid have been reported: antifibrotic activity, protection of neurons against insults, suppression of UVB-induced alterations to human keratinocytes,



inhibition of bone metastasis from breast carcinoma<sup>[20]</sup>.

Salvianolic acid K is reported in the literature for its antioxidant<sup>[21]</sup> and aldose reductase inhibitory activities<sup>[22]</sup>.

Among flavones, cynaroside (luteolin-7-O-β-D-glucopyranoside) prevents H<sub>2</sub>O<sub>2</sub>-induced apoptosis in H9c2<sup>[23]</sup> and SH-SY5Y<sup>[24]</sup> cell lines by reducing the endogenous production of ROS. Furthermore, cynaroside is very effective against Gram-negative bacteria<sup>[25]</sup> and showed significant antipsoriatic activity using photodermatitis model<sup>[26]</sup>.

Isocutellarein derivatives have also been described for its important beneficial activities: antioxidant<sup>[27]</sup> antinociceptive and anti-inflammatory<sup>[28]</sup> and significant inhibitory activity against osteoclast differentiation<sup>[29]</sup>.

Spices are widely investigated for their essential oils content in term of chemical composition and biological activity. Nevertheless, spices are also abundant sources of polyphenols which have interesting biological properties as recently reported also by Kindl, et al (2015)<sup>[30]</sup> that report antioxidant and acetylcholinesterase inhibitory activities of the ethanolic extracts of six selected *Thymus* species growing in Croatia. Consumption of spices has been implicated in the prevention of cardiovascular diseases, carcinogenesis, and inflammation mainly due to the presence of polyphenols.

## References

- Meskin, M.S., Bidlack, W.R., Davies, A.J., et al. Phytochemicals in Nutrition and Health. (2002) CRC Press.
- Hossain, M.B., Brunton, N.P., Barry-Ryan, C., et al. Antioxidant activity of spice extract and phenolics in comparison to synthetic antioxidants. (2008) *Rasayan J Chem* 1(4): 751-756.
- Saxena, M., Saxena, J., Pradhan, A. Flavonoids and phenolic acids as antioxidant in plants and human health. (2012) *Int J Pharm Sci* 16(2): 130-134.
- De-Martino, L., Bruno, M., Formisano, C., et al. Chemical composition and antimicrobial activity of the essential oils from two species of *Thymus* growing wild in southern Italy. (2009) *Molecules* 14(11): 4614-4624.
- Scherrer, A.M., Motti, R., Weckerle, C.S. Traditional plant use in the areas of Monte Vesole and Ascea, Cilento National Park (Campania, Southern Italy). (2005) *J Ethnopharmacol* 97(1): 129-143.
- Vladimir-Knežević, S., Kosalec, I., Babac, M., et al. Antimicrobial activity of *Thymus longicaulis* C. Presl essential oil against respiratory pathogens. (2012) *Cent Eur J Biol* 7(6): 1109-1115.
- Sarikurkcu, C., Sabih-Ozer, M., Eskici, M., et al. Essential oil composition and antioxidant activity of *Thymus longicaulis* C. Presl subsp. *longicaulis* var. *longicaulis*. (2010) *Food Chem Toxicol* 48(7): 1801-1805.
- Scognamiglio, M., D'Abrosca, B., Esposito, A., et al. Chemical Composition and Seasonality of Aromatic Mediterranean Plant Species by NMR-Based Metabolomics. (2015) *J Anal Methods Chem* 258570.
- Piccolella, S., Fiorentino, A., Pacifico, S., et al. Antioxidant properties of sour cherries (*Prunus cerasus* L): role of colorless phytochemicals from the methanolic extract of ripe fruits. (2008) *J Agric Food Chem* 56(6): 1928-1935.
- Cui, B., Nakamura, M., Kinjo, J., et al. Chemical Constituents of *Astragali Semen*. (1993) *Chem Pharm Bull* 41(1): 178-182.
- Cefarelli, G., D'Abrosca, B., Fiorentino, A., et al. Free-radical-scavenging and antioxidant activities of secondary metabolites from reddened cv. Annurca apple fruits. (2006) *J Agric Food Chem* 4(3): 803-809.
- Tezuka, Y., Kasimu, R., Xin, Li. Constituents of roots of *Salvia deserti*. (1998) *Chem Pharm Bull* 46 107-112.
- Zhou, L.Y., Liu, H-Y., Xie, B.B., et al. Two new glycosides from *Sanicula lamelligera*. Z. (2006) *Naturforsch* 61b: 607-610.
- D'Abrosca, B., Pacifico, S., Scognamiglio, M., et al. A new acylated flavone glycoside with antioxidant and radical scavenging activities from *Teucrium polium* leaves. (2013) *Nat Prod Res* 27(4-5): 356-363.
- Rigano, D., Formisano, C., Basile, A., et al. Antibacterial activity of flavonoids and phenylpropanoids from *Marrubium globosum* ssp. *libanoticum*. (2007) *Phytother Res* 21(4): 395-397.
- Iida, N., Inatomi, Y., Murata, H., et al. A new flavone xyloside and two new flavan-3-ol glucosides from *Juniperus communis* var. *depressa*. (2007) *Chem Biodivers* 4(1): 32-42.
- Schliemann, W., Schneider, B., Wray, V., et al. Flavonols and an indole alkaloid skeleton bearing identical acylated glycosidic groups from yellow petals of *Papaver nudicaule*. (2006) *Phytochemistry* 67(2): 191-201.
- Kizu, H., Shimana, H., Tomimori, T. Studies on the constituents of *Clematis* species. (1995) *Chem Pharm Bull* 43(12): 2187-2194.
- Petersen, M., Simmonds, M.J. Rosmarinic acid. (2003) *Phytochemistry* 62(2): 121-125.
- Bulgakov, V.P., Inyushkina, Y.V., Fedoreyev, S.A. Rosmarinic acid and its derivatives: biotechnology and applications. (2012) *Crit Rev Biotechnol* 32(3): 203-217.
- Bors, W., Michel, C., Stettmaier, K., et al. Antioxidant mechanisms of polyphenolic caffeic acid oligomers, constituents of *Salvia officinalis*. (2004) *Biol Res* 37(2): 301-311.
- Kasimu, R., Tanaka, K., Tezuka, Y., et al. Comparative study of seventeen *Salvia* plants: Aldose reductase inhibitory activity of water and MeOH extracts and liquid chromatography-mass spectrometry (LC-MS) analysis of water extracts. (1998) *Chem Pharm Bull* 46(3): 500-504.
- Sun, X., Sun, G.B., Wang, M., et al. Protective effects of cynaroside against H<sub>2</sub>O<sub>2</sub>-induced apoptosis in H9c2 cardiomyoblasts. (2011) *J Cell Biochem* 112(8): 2019-2029.
- Zhang, Y.C., Gan, F.F., Shelar, S.B., et al. Antioxidant and Nrf2 inducing activities of luteolin, a flavonoid constituent in *Ixeris sonchifolia* Hance, provide neuroprotective effects against ischemia-induced cellular injury. (2013) *Food Chem Toxicol* 59: 272-280.
- Žemlička, L., Fodran, P., Lukeš, V., et al. Physicochemical and biological properties of luteolin-7-O-β-D-glucoside (cynaroside) isolated from *Anthriscus sylvestris* (L.) Hoffm. (2014) *Monatsh Chem* 145(8): 1307-1318.
- Vijayalakshmi, A., Geetha, M. Anti-psoriatic activity of flavonoids from *Cassia tora* leaves using the rat ultraviolet B ray photodermatitis model. (2014) *Rev Bras Farmacogn* 24(3): 322-329.
- Yang, H., Protiva, P., Cui, B., et al. New bioactive polyphenols from *Theobroma grandiflorum* ("cupuaçu"). (2003) *J Nat Prod* 66(11): 1501-1504.
- Küpeli, E.E., Sahin, F.P., Yeşilada, E., et al. In vivo anti-inflammatory and antinociceptive activity evaluation of phenolic compounds from *Sideritis stricta*. (2007) *Z Naturforsch* 62(7-8): 519-525.
- Yoon, K.D., Jeong, D.G., Hwang, Y.H., et al. Inhibitors of osteoclast differentiation from *Cephalotaxus koreana*. (2007) *J Nat Prod* 70(12): 2029-2032.
- Kindl, M., Blažeković, B., Bucar, F., et al. Antioxidant and Anticholinesterase Potential of Six *Thymus* Species. (2015) *Evid Based Complement Alternat Med* 2015: 403950.